



UNIVERSITI PUTRA MALAYSIA

**DISEASE CAUSED BY PHYTIUM SPP. ON SELECTED CROPS
GROWN UNDER HYDROPONIC CULTURE SYSTEM AND THEIR
CONTROL**

CHIN CHIEW LAN

FP 1988 7

DISEASES CAUSED BY PYTHIUM SPP. ON SELECTED
CROPS GROWN UNDER HYDROPONIC CULTURE SYSTEM
AND THEIR CONTROL

BY


CHIN CHIEW LAN

A thesis submitted in partial fulfilment of the
requirements for the degree of Master of Agricultural Science
in the Faculty of Agriculture,
Universiti Pertanian Malaysia

September 1988



It is hereby certified that we have read this thesis entitled 'Diseases Caused by Pythium Spp. on Selected Crops Grown under Hydroponic Culture System and their Control' by Chin Chiew Lan, and in our opinion it is satisfactory in terms of the scope, quality, and presentation as partial fulfilment of the requirements for the degree of Master of Agricultural Science



.....
SULAIMAN M. YASSIN, Ph.D.
Professor/Dean of Graduate Studies
Universiti Pertanian Malaysia
(Chairman Board of Examiners)



.....
WEN-HSIUNG KO, Ph.D.
Professor
Department of Plant Pathology
University of Hawaii, Beaumont
U.S.A.
(External Examiner)



.....
ZAINAL ABIDIN MIOR AHMAD
Faculty of Agriculture
Universiti Pertanian Malaysia
(Internal Examiner)



.....
LIM TONG KWEE, Ph.D.
Associate Professor
Faculty of Agriculture
Universiti Pertanian Malaysia
(Internal Examiner/Supervisor)

This thesis was submitted to the Senate of Universiti Pertanian Malaysia and was accepted as partial fulfilment of the requirements for the degree of Master of Agricultural Science.

Date: 10 NOV 1975


.....
Dean of Graduate School

Prof. Dr. Sulaiman M. Yassin
Dean of Graduate School
Universiti Pertanian Malaysia
43400 Serdang
SELANGOR MALAYSIA.

ACKNOWLEDGEMENTS

The author wishes to thank in particular her supervisor, Dr. Lim Tong Kwee and co-supervisor, Mr. Abdul Khalip Raffar for their invaluable guidance, advice and constructive criticisms throughout the duration of the project.

Appreciations are also extended to the academic staff of Universiti Pertanian Malaysia especially Dr. Raja Muhammad Harun, Dr. Lim Eng Seong and Dr. Surjit Singh for their useful advice; staff of Hydroponic Unit of Universiti Pertanian Malaysia especially Mr. Abdul Ghani Hashim, Mr. Abdul Aziz Abdul Rahman and Mr. Mohd. Yunus Abdul Wahab and laboratory staff of the Department of Plant Protection, Universiti Pertanian Malaysia for their kind assistance and cooperation, Mr. Zainal Abidin Mior Ahmad for translating the abstract, and Miss Low Kwee Choon for taking the pains and extra hours typing the manuscript.

The author also wishes to thank Universiti Pertanian Malaysia for providing assistantship to the author in her first year and SEARCA for sponsorship of her second year of post-graduate studies in Universiti Pertanian Malaysia.

Finally, the author owes a great deal to her family and fiance Pou Hai for their love, patience and encouragement.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	ii
TABLE OF CONTENTS	iii
LIST OF TABLES	vii
LIST OF FIGURES	ix
LIST OF PLATES	xi
ABSTRACT	xiv
ABSTRAK	xix
 CHAPTER 1 INTRODUCTION	 1
 CHAPTER 2 CULTURAL AND MORPHOLOGICAL STUDIES	
INTRODUCTION	6
LITERATURE REVIEW	6
Taxonomy of <u>Pythium</u>	6
Identification of <u>Pythium</u> spp.	8
Colony Pattern	8
Appressoria	9
Sporangia and hyphal swellings	9
Antheridia, oogonia, oospores and heterothallism	10
Temperature relationship	13
MATERIALS AND METHODS	13
Isolation	13
Effect of media and temperature on growth rate and colony morphology	19
Production of asexual morphological characters	21



	Page
Production of sexual morphological characters	22
RESULTS	23
Effect of media and temperature on growth rate and colony morphology	23
Linear mycelial growth	23
Growth rate	29
Colony morphology	37
Production of asexual morphological characters	38
Appressoria	38
Sporangia	39
Hyphal swellings	45
Production of sexual morphological characters	45
Oogonia, antheridia and oospores	45
DISCUSSION	52
CHAPTER 3 PATHOGENICITY STUDIES	
INTRODUCTION	63
LITERATURE REVIEW	64
<u>Pythium</u> in soil versus <u>Pythium</u> in hydroponic culture system	64
MATERIALS AND METHODS	67
Pathogenicity of <u>Pythium</u> isolates on cucumber, muskmelon and tomato	67
Differential response of <u>Pythium</u> isolates on cucumber seedlings	68

	Page
Pathogenicity of <u>Pythium</u> isolates on cucumber and muskmelon at different crop growth stages	70
Sources and inoculum level of <u>Pythium</u> in the Universiti Pertanian Malaysia commercial hydroponic system	74
RESULTS	76
Pathogenicity of <u>Pythium</u> isolates on cucumber, muskmelon and tomato	76
Differential response of <u>Pythium</u> isolates on cucumber seedlings	76
Pathogenicity of <u>Pythium</u> isolates on cucumber and muskmelon at different crop growth stages	80
Sources and inoculum level of <u>Pythium</u> in the Universiti Pertanian Malaysia Commercial Hydroponic System.	89
DISCUSSION	92
CHAPTER 4 CONTROL STUDIES	
INTRODUCTION	101
LITERATURE REVIEW	101
Control of <u>Pythium</u> diseases in hydroponic culture system	101
MATERIALS AND METHODS	104
<u>In-vitro</u> studies	104
Effect on linear mycelial growth	106
<u>In-vivo</u> studies	106
Relationship of inoculum concentration and disease level	106
Efficacy of chemicals on pre-emergent damping-off of cucumber seedlings	106

	Page
Efficacy of chemicals on post-emergent damping-off of cucumber seedlings	108
RESULTS	110
<u>In-vitro</u> studies	110
Effect on linear mycelial growth	110
<u>In-vivo</u> studies	113
Relationship of inoculum concentration and disease level	113
Efficacy of chemicals on pre-emergent damping-off of cucumber seedlings	113
Efficacy of chemicals on post-emergent damping-off of cucumber seedlings	115
DISCUSSION	118
CHAPTER 5 SUMMARY AND CONCLUSION	126
BIBLIOGRAPHY	131
APPENDICES	137

LIST OF TABLES

TABLE NO.	DESCRIPTION	PAGE
1	Isolates of <u>Pythium</u> used in the studies	20
2	Polynomials expressing the effect of temperature (t) upon the linear growth diameter (Y) of six <u>Pythium</u> isolates grown in: (a) CMA, (b) VJA and (c) HNA	27
3	Comparison among isolates in the study of the effect of media and temperature on mycelial linear growth of <u>Pythium</u> isolates	30
4	Comparison among media in the study of the effect of media and temperature on mycelial linear growth of <u>Pythium</u> isolates	31
5	Polynomials expressing the effect of temperature (t) upon the growth rate (Z) of six <u>Pythium</u> isolates grown in: (a) CMA, (b) VJA and (c) HNA	35
6	Measurement of oogonia and oospores of <u>Pythium</u> isolates on cornmeal broth (CMB) after 4 days of incubation at 28 ± 1 C	49
7	Effect of inoculation with six <u>Pythium</u> isolates on the germination of cucumber, muskmelon, and tomato seed	77
8	Pathogenicity of <u>Pythium</u> isolates on cucumber seedlings	78
9	Height of cucumber plants inoculated with <u>Pythium</u> isolate Py 15 at different crop growth stage	86
10	Influence of inoculating cucumber plants with <u>Pythium</u> isolate Py 15 at different crop growth stages on the yield response at 60 days after sowing	87



		PAGE
11	Effect of inoculating cucumber plants with <u>Pythium</u> isolate Py 15 at different crop growth stages on the root at 60 days after sowing	88
12	Effect of inoculating muskmelon plant with <u>Pythium</u> isolate Py 5 at different crop growth stages on the yield response at 70 days after sowing	90
13	Fungicides used in the toxicity studies of <u>Pythium</u> spp. isolated from crops grown under hydroponic culture system	105
14	Fungicides and rates used in pre-emergent damping-off evaluation	107
15	Fungicides and rates used in post-emergent damping-off evaluation	109
16	Regression equation expressing the toxicity of chemicals on <u>Pythium</u> isolates (Py 9, 15, 23)	111
17	Toxicity of chemicals on mycelial growth of <u>Pythium</u> isolates (Py 9, 15, 23) at 18 hours of incubation	112
18	Effect of fungicides at various concentrations applied as seed treatment in the control of pre-emergent damping-off of cucumber seed caused by <u>Pythium myriotylum</u> (isolate Py 15)	116
19	Effect of fungicides applied to the nutrient solution in the evaluation of post-emergent damping-off caused by <u>Pythium myriotylum</u> (isolate Py 15)	117



LIST OF FIGURES

FIGURE NO.	DESCRIPTION	PAGE
1	Layout of the two-phase substrate culture system. (After Lim and Wan, 1984)	2
2	Effect of temperature on mycelial linear growth of <u>Pythium</u> isolates on cornmeal agar (CMA) after 18 hours of incubation	24
3	Effect of temperature on mycelial linear growth of <u>Pythium</u> isolates on V-8 juice agar (VJA) after 18 hours of incubation	25
4	Effect of temperature on mycelial linear growth of <u>Pythium</u> isolates on Hyponica nutrient agar (HNA) after 18 hours of incubation.	26
5	Mean hourly growth of six <u>Pythium</u> isolates on cornmeal agar (CMA) in relation to temperature	32
6	Mean hourly growth of six <u>Pythium</u> isolates on V-8 juice agar (VJA) in relation to temperature	33
7	Mean hourly growth of six <u>Pythium</u> isolates on Hyponica nutrient agar (HNA) in relation to temperature	34
8	Appressoria as seen from the underside of petri dish culture. a. Appressoria formed in clusters consisting of globose or knob-like terminations. b, c. Appressoria in the early stage of formation. d,e,f,g,h. Sickle-, club- or sausage-shaped or subglobose forms of appressoria. i. swollen digitate sporangia.	39



		PAGE
9	a. Encysted zoospore. b,c,d. Germinating zoospores with one, two, three germ tubes respectively.	43
10	a,b. Terminal and intercalary oogonia c,d. Diclinous antheridia; branched antheridia stalk. e,f. Clavate; crook-necked or arched antheridia	48
11	a. Crop phenology of cucumber (Cultivar Burpless Green King)	71
	b. Crop phenology of muskmelon (Cultivar Birdie)	73
12	The percentage frequency of <u>Pythium</u> isolation from root tissue of muskmelon plants grown in the commercial hydroponic culture system in relation to the crop phenology.	93
13	The relative level of <u>Pythium</u> in the nutrient solution in relation to the crop phenology of muskmelon grown in the commercial hydroponic culture system.	94
14	Linear relationship between percentage pre-emergent kill of cucumber seedlings (probit scale) and concentration of <u>Pythium myriotylum</u> (Py 15) in the nutrient solution (logarithmic scale)	114

LIST OF PLATES

PLATE NO.	DESCRIPTION	PAGE
I	a. Interveinal chlorosis of muskmelon leaves at the fruit development stage.	14
	b. Highly decayed root mat of muskmelon plant showing absence of root hairs	15
II	Root mat of muskmelon grown in FIMA's Fraser's Hill hydroponic unit showing brown discoloured patch	16
III	a. Wilted cucumber plants growing in the commercial hydroponic culture system (Serdang)	17
	b. 'Damping-off' of cucumber seedlings in the nursery stage	18
IV	a. Sausage - or club-shaped appressoria of <u>Pythium myriotylum</u> (Py 36). Bar: 1 cm = 7.69 μ m	40
	b. Appressoria of <u>P. myriotylum</u> (Py 36) formed singly or connected to each other in a chain	
V	Lobulate sporangia of <u>Pythium myriotylum</u> (Py 9)	41
VI	Lobulate sporangia of <u>Pythium myriotylum</u> (Py 9) arranged in close lateral clusters	42
VII	Encysted zoospores of <u>Pythium myriotylum</u> (Py 15) germinating by a single germ tube (arrowed) Bar: 1 cm = 5.2 μ m	44
VIII	Hyphal swelling of <u>Pythium splendens</u> (Py 23) with dark and densely granulated protoplasm. Bar: 1 cm = 14.8 μ m	46
IX	Hyphal swelling of <u>Pythium splendens</u> (Py 23) with a large reniform oil globule (arrowed) Bar: 1 cm = 6 μ m	47



X	Diclinous antheridia of <u>Pythium myriotylum</u> (Py36) with a branched antheridial stalk (arrowed) a. Bar: 1 cm = 9.0 μ m b. Bar: 1 cm = 6.7 μ m	50
XI	Clavate, crook neck or arched antheridia of <u>Pythium myriotylum</u> (Py 15) (arrowed). Bar: 1 cm = 5.7 μ m	51
XII	Oogonium of <u>Pythium myriotylum</u> (Py 5) with smooth wall being partially enveloped by a cluster of branched antheridia. Bar: 1 cm = 2.9 μ m	53
XIII	Aplerotic oospores of <u>Pythium myriotylum</u> (Py 9) with reserve globule in the protoplasm (arrowed) Bar: 1 cm = 5.6 μ m	54
XIV	Oospore of <u>Pythium myriotylum</u> (Py 9) granulated protoplasm. Bar: 1 cm = 6.6 μ m	55
XV	Slightly inflated filamentous dendroid hyphae of <u>Pythium</u> isolate Py 19	61
XVI	Cucumber seedlings at five days after sowing showing wilted symptom at one day after inoculation with <u>Pythium splendens</u> (Py 23)	79
XVII	a. Cucumber seedlings showing damping-off when inoculated with <u>Pythium myriotylum</u> (Py 15) at 2-3 days after sowing b. The check plot showing healthy cucumber seedlings	81
XVIII	Cucumber seedlings showing damping-off near the basal portion with water-soaked lesions which frequently advance up the stem for 2 - 3 cm before drying-up and falling over	82

	PAGE
XIX	Cucumber plants inoculated with <u>Pythium myriotylum</u> (Py 15) at 30 days after sowing, growing luxuriantly with thick large leaves in non-aerated nutrient solution in the glass house 84
XX	Typical white healthy root mat of cucumber plant inoculated with <u>Pythium myriotylum</u> (Py 15) at 14, 30 or 40 days after sowing 85
XXI	Oospores of <u>Pythium myriotylum</u> present in root tissue of muskmelon Bar : 1 cm = 19 um 91

ABSTRACT

An abstract of the thesis presented to the Senate of Universiti Pertanian Malaysia in partial fulfilment of the requirements for the degree of Master of Agricultural Science.

DISEASES CAUSED BY PYTHIUM SPP. ON SELECTED CROPS GROWN UNDER HYDROPONIC CULTURE SYSTEM AND THEIR CONTROL

by

Chin Chiew Lan

1988

Supervisor : Associate Prof. Dr. Lim Tong Kwee
Co-supervisor : Associate Prof. Khalip Abdul Raffar
Faculty : Agriculture

Large scale hydroponic crop production is a recent entity in Malaysia with University of Agriculture Malaysia (UPM) playing a lead role in research and development. Since its inception in 1982, pre- and post-emergent damping-off in cucumber, muskmelon and tomato seedlings had been frequently observed. Pythium spp. were also isolated from roots of wilted mature cucumber, muskmelon and tomato plants at UPM hydroponic, at Malaysia Agricultural Research and Development Institute (MARDI) and at FIMA hydroponic in Fraser's Hill. The mature muskmelon plants exhibited drastic interveinal chlorosis and yellowing of young and old leaves at the onset of

fruiting and was accompanied by general necrosis of the root system resulting in subsequent plant death. Isolation of Pythium implicate them to be the causal agent. Since little is known about their role and control in the hydroponic system in Malaysia, studies were undertaken to characterize and identify the Pythium spp., to establish their pathogenicity, to determine the sources of inoculum, as well as to investigate the in-vitro and in-vivo efficacy of some fungicides in their control.

Through cultural and morphological studies, three Pythium spp. namely P. myriotylum, P. splendens and a non-sporulating Pythium sp. were identified from six isolates obtained.

P. myriotylum was isolated from muskmelon, cucumber and tomato grown in the hydroponic culture system in UPM, Serdang. P. myriotylum was characterized by its optimum temperature for growth at 35°C, and its ability to grow at 40°C and above. It produced numerous clavate, knob-like, or sickle-shaped appressoria. Inflated, digitate sporangia which discharged zoospores were observed in grass-blade water culture. It formed abundant oogonia with less than 10 diclinous antheridia in single cultures. Oospores were highly aplerotic with diameter ranging from 17.65 ± 0.72 to 21.15 ± 1.00 μm .

P. splendens was isolated from muskmelon in FIMA hydroponic set-up at Fraser's Hill. It produced numerous large spherical terminal hyphal swellings in culture which measured 40.4 ± 1.04 μm in diameter. It did not form oogonia nor oospore in single

cultures. The optimum and maximum temperature for growth was 30°C and 35°C respectively.

The unidentified Pythium isolated from tomato plant grown in the UPM hydroponic unit at Genting Highlands, produced slightly inflated, filamentous, dendroid hyphae which did not discharge zoospores. It neither formed hyphal swellings, appressoria nor any sexual reproductive structures. It grew optimumly at 30°C, and maximumly at 40°C. Protoplasmic streaming in young hyphae were observed.

The predominant Pythium sp. isolated in UPM hydroponic unit at Serdang, was P. myriotylum, and pathogenicity studies showed that it caused pre-emergent damping-off of cucumber, muskmelon and tomato; and post-emergent death of cucumber and muskmelon seedlings younger than 7 days after sowing (DAS). Cucumber and muskmelon plants older than 7 DAS appeared to be tolerant to P. myriotylum infection.

P. splendens isolated from the FIMA hydroponic set-up at Fraser's Hill and the unidentified Pythium sp. from UPM hydroponic unit at Genting Highlands, were pathogenic on all three crops namely cucumber, muskmelon and tomato at the pre-emergent stage.

All six isolates of Pythium viz P. myriotylum (4 isolates), P. splendens (1 isolate) and Pythium sp. (1 isolate) exhibited differential pathogenic response and isolates of the same species (P. myriotylum) varied in their pathogenic response. The pathogenicity studies further revealed that interveinal yellowing,

root death and subsequent plant mortality of mature muskmelon was not related to P. myriotylum infection. This was further confirmed by other University scientists who ascertained that the above symptoms were more the result of manganese and iron deficiency.

The inoculum source and population studies showed that P. myriotylum was present in the hydroponic nutrient solution and roots of both mature healthy as well as chlorotic muskmelon plant. In addition, the fluctuations in population level of Pythium were related to the phenology of the crop. P. myriotylum was also found to be present throughout the whole nutrient solution in the system.

In-vitro chemical efficacy studies indicated that the fungitoxicity varied according to Pythium sp. For P. myriotylum, etridiazole was the most toxic, and metalaxyl was the next most potent. Next in efficacy ranked propamocarb hydrochloride, oxadixyl plus mancozeb, copper hydroxide and phosetyl-Al in descending effectiveness. For P. splendens, on the other hand, metalaxyl was the most toxic; followed by etridiazole, oxadixyl plus mancozeb, copper hydroxide, propamocarb hydrochloride and phosetyl-Al in descending efficacy.

None of the four fungicides namely Copper hydroxide, metalaxyl, etridiazole and propamocarb hydrochloride outperformed each other significantly in the pre-emergent efficacy study. When applied into the nutrient solution as post-emergent treatment, metalaxyl was the most effective, followed by copper hydroxide, etridiazole and

oxadixyl plus mancozeb which did not differ significantly in the control of the disease. Phosetyl-Al and propamocarb hydrochloride showed negligible effect.

ABSTRAK

Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi sebahagian daripada syarat-syarat untuk Ijazah Master Sains Pertanian.

PENYAKIT-PENYAKIT OLEH PYTHIUM SPP. PADA TANAMAN- TANAMAN TERPILIH DITANAM DI BAWAH SISTEM KULTUR HIDROPONIK DAN KAWALANNYA

Oleh

Chin Chiew Lan

1988

Penyelia : Prof. Madya Dr. Lim Tong Kwee

Penyelia Bersama : Prof. Madya Khalip Abdul Raffar

Fakulti : Pertanian

Pengeluaran secara besar-besaran tanaman hidroponik adalah satu kewujudan baru di Malaysia dengan Universiti Pertanian Malaysia (UPM) memainkan peranan utama di dalam penyelidikan dan perkembangannya. Sejak bermulanya dari tahun 1982, penyakit melecur sebelum dan selepas cambah pada tanaman anak benih timun, tembikai wangi dan tomato didapati kerap berlaku. Pythium spp. telah juga dipencilkan dari akar-akar tanaman matang tembikai wangi, timun dan tomato yang layu di kawasan tanaman hidroponik, UPM, Institut Penyelidikan dan Kemajuan Pertanian Malaysia (MARDI) dan pada tanaman hidroponik FIMA di Fraser's Hill. Tembikai wangi matang

menunjukkan tanda-tanda klorosis di antara urat-urat daun yang amat ketara dan penguningan daun-daun muda dan tua pada peringkat permulaan membuah dan diikuti oleh nekrosis am sistem akar yang menyebabkan kematian tanaman. Pencilan Pythium mengaitkannya sebagai agen penyebab penyakit berkenaan. Oleh kerana hanya sedikit diketahui tentang peranan dan kawalannya di dalam sistem tanaman hidroponik di Malaysia, kajian-kajian telah dijalankan untuk perbandingan ciri-ciri dan pengenalan Pythium spp., menguji kepatogenan, menentukan punca-punca inokulum, di samping mengkaji keberkesanan beberapa racun kulat secara in-vitro dan in-vivo bagi kawalannya.

Melalui kajian-kajian kultur dan morfologi, tiga Pythium spp. iaitu P. myriotylum, P. splendens dan satu Pythium sp. yang tidak mengeluarkan spora, telah dicam daripada enam pencilan-pencilan yang didapati.

P. myriotylum telah dipencil daripada tembikai wangi, timun dan tomato yang ditanam di bawah sistem kultur hidroponik di UPM, Serdang. P. myriotylum telah dikenalpasti melalui ciri-ciri suhu optimum pada 35°C bagi pertumbuhannya dan keupayaan tumbuh pada 40°C ke atas. Ia mengeluarkan dengan banyak appressoria yang berbentuk klavat, seperti gembul atau sabit. Sporangium digitat yang membesar yang mengeluarkan zoospora didapati di dalam sistem kultur air bersama lai rumput. Oogonium telah didapati dengan banyak dan mempunyai kurang dari sepuluh anteridia diklinus dalam kultur

tunggal. Oospora adalah amat aplerotik dan mempunyai garis pusat berjulat antara 17.65 ± 0.72 hingga 21.15 ± 1.00 μm .

P. splendens telah dipencil daripada tembikai wangi pada Unit Hidroponik FIMA di Fraser's Hill. Ia mengeluarkan dengan banyak bengkakan hifa yang besar dan bulat di dalam kultur yang mempunyai garis pusat berukuran 40.4 ± 1.04 μm . Ia tidak membentuk oogonium atau oospora di dalam kultur tunggal. Suhu optimum untuk pertumbuhan ialah 30°C manakala suhu maksimum ialah 35°C .

Pythium yang tidak dapat dikenalpasti telah dipencil daripada tomato yang ditanam di Unit Hidroponik UPM di Genting Highlands, yang mengeluarkan hifa dendroid, berfilamen dan hanya membesar sedikit dan tidak mengeluarkan zoospora. Ia tidak membentuk bengkakan hifa, appressorium atau apa-apa struktur pembiakan seks. Ia mencapai pertumbuhan optimum pada 30°C , manakala pertumbuhan maksima pada 40°C . Aliran protoplasma di dalam hifa muda telah dilihat.

Pythium sp. utama yang telah dipencilkan di UPM, Serdang ialah P. myriotylum, dan ujian kepatogenan menunjukkan yang ia menyebabkan penyakit melecur sebelum dan selepas cambah tanaman timun, tembikai wangi dan tomato; dan juga kematian selepas cambah anak benih timun dan tembikai wangi yang berusia muda iaitu tidak melebihi 7 hari selepas semaian. Tanaman timun dan tembikai wangi yang lebih tua daripada 7 hari selepas semaian didapati tahan terhadap jangkitan P. myriotylum.

P. splendens yang dipencil daripada Unit Hidroponik FIMA di Fraser's Hill dan Pythium sp. yang tidak dikenalpasti daripada Unit Hidroponik UPM, Genting Highlands didapati adalah merupakan patogen kepada ketiga-tiga jenis tanaman iaitu timun, tembikai wangi dan tomato pada peringkat sebelum cambah.

Kesemua 6 pencilan-pencilan Pythium iaitu P. myriotylum (4 pencilan), P. splendens (1 pencilan) dan Pythium sp. (1 pencilan) menunjukkan perbezaan respon kepatogenan dan pencilan yang sama daripada spesies (P. myriotylum) menunjukkan respon kepatogenan yang berlainan. Kajian kepatogenan juga menunjukkan bahawa kekuningan di antara urat-urat daun, kematian akar dan kematian pada tembikai wangi matang adalah tidak berkaitan dengan jangkitan P. myriotylum. Ini telah turut disahkan oleh lain-lain saintis universiti yang telah menunjukkan bahawa simptom-simptom di atas adalah lebih merupakan akibat kekurangan mangan dan besi.

Punca inokulum dan kajian populasi menunjukkan bahawa P. myriotylum terdapat di dalam larutan pemakanan hidroponik dan akar-akar kedua-dua tanaman tembikai wangi matang yang sihat dan juga yang menunjukkan simptom klorosis. Di samping itu, turun naik paras populasi Pythium didapati berhubung dengan fenologi tumbuhan tersebut. P. myriotylum telah juga didapati pada seluruh kandungan larutan pemakanan di dalam sistem berkenaan.

Kajian keberkesanan kimia in-vitro menunjukkan bahawa keracunan racun kulat adalah berlainan mengikut Pythium sp. Bagi P. myriotylum, etridiazole adalah paling beracun, diikuti berikutnya